

An Entomological Surveillance System Based on Open Spatial Information for Participative Dengue Control

LEDA REGIS¹, WAYNER VIEIRA DE SOUZA², ANDRÉ FREIRE FURTADO¹, CLAUDIO DUARTE DA FONSECA³, JOSÉ CONSTANTINO SILVEIRA Jr², PAULO JUSTINIANO RIBEIRO Jr⁴, MARIA ALICE VARJAL DE MELO-SANTOS¹, MARÍLIA SA CARVALHO⁵, ANTONIO MIGUEL VIEIRA MONTEIRO⁶

¹Departamento de Entomologia, Centro de Pesquisas Aggeu Magalhães, Fundação Oswaldo Cruz, Cx Postal 7472 Cidade Universitária, Recife 50670-420 Recife, PE, Brasil.

²Departamento de Saúde Coletiva, Centro de Pesquisas Aggeu Magalhães, Fundação Oswaldo Cruz, Av. Prof. Moraes Rego s/n, 50670-420, Recife, PE, Brasil.

³Secretaria Estadual de Saúde de Pernambuco, Praça Oswaldo Cruz s/n Boa Vista, 50050-210 Recife, PE, Brasil

⁴Laboratório de Estatística e Geoinformação, Departamento de Estatística, Universidade Federal do Paraná-UFPR, Curitiba, PR, Brasil

⁵Escola Nacional de Saúde Pública –ENSP, Fundação Oswaldo Cruz, Rio de Janeiro, RJ, Brasil

⁶Divisão de Processamento de Imagens & Programa Espaço e Sociedade, Instituto Nacional de Pesquisas Espaciais-INPE, São José dos Campos, SP, Brasil

Key words: dengue vector, entomological surveillance, ovitraps, GIS, vector control

An entomological surveillance system for dengue control

Biological Sciences

Lêda Regis, CPqAM-Fiocruz, Cx Postal 7472 CEP 50670-420, Recife, PE, Brasil. Fax: 55-81-21012516. Tel.: 55-81-21012552. E-mail: leda@cpqam.fiocruz.br

ABSTRACT

Aedes aegypti is a very efficient disseminator of human pathogens as a result of evolutionary adaptations to frequent haematophagy and almost exclusively in humans as well as to the colonization of countless types of habitats, associated with biotic, climate and cultural factors that favors the proliferation of this mosquito in urban ecosystems. Studies using sensitive methods of monitoring demonstrate that the methods of surveillance used in the Brazilian program do not show the high intensities of the infestation of cities by this vector. To enlarge the capacity of the health sector new tools are necessary to the practice of surveillance, incorporating aspects of the vector, of the place and of the local human population. We describe here the SMCP-*Aedes* - Monitoring System and Population Control of *Aedes aegypti*, which goal is the instrumentalization of entomological surveillance as a basis for epidemiological surveillance of dengue. The SMCP-*Aedes* is uphold in the space technology information, supported by the intensive use of the web and free software to collect, store, analyze and disseminate information on the spatial-temporal distribution of the estimated density for the population of *Aedes*, based on data systematically collected with the use of ovitraps. Planned control interventions, intensified where and when indicated by the entomological surveillance, are agreed with the communities, relying on the permanent social mobilization.

INTRODUCTION

According to the World Health Organization, over 40% of the world's population is at risk of acquiring dengue and there may be 50 million of new dengue infections every year. Innate characteristics of the dengue virus (DENV), which exist in four serotypes and can cause up to four infections per person, and of its main vector, *Aedes aegypti*, considered an extremely

effective disseminator of human pathogens, account for the expansion of the geographic distribution and, consequently, for the dramatic growth of the global burden of dengue in the last decades. *Ae. aegypti* is a highly anthropophilic species, that feeds almost exclusively on humans (Scott et al 1993, 2000, Harrington et al 2001, Ponllawat and Harrington 2005). According to Harrington et al. (2001), the unique isoleucine concentration on human blood is associated with *Ae. aegypti*'s unusual propensity to preferentially and frequently feed on humans – a behavior that increases this mosquito's fitness and thus is an underlying reason for its intimate contact with human hosts. The authors highlight that “from a public health perspective, selection for frequent human contact and an age-dependent increase in feeding frequency magnify the potential for transmission of pathogens by older potentially infected mosquitoes”. Upon studying the blood meal frequency of *Ae. aegypti* field populations in Thailand, Scott et al. (2000) estimated that, on average, a female takes 0.76 human blood meal per day. These biological features of the vector have high epidemiological significance, and explain why *Ae. aegypti*'s territory is the human house, from which it seldom leaves.

Evolution driven adaptations to unstable habitats are also important factors to understand the role of this species as an efficient vector of pathogens: *Ae. aegypti* is an opportunistic container breeder. This mosquito is able to colonize all types of water holding containers; presents a fantastic resistance of its eggs chorion and the “skip oviposition behavior” of females (i.e. the spread of eggs from the same batch on different containers); yonder the ability of larvae and pupae to dive deeper and for longer periods than other mosquito species (Koenraadt and Harrington, 2008). It is crucial to consider these biological aspects and the consequences of their combination with environmental and climatic factors, and with anthropogenic habits that favor mosquito proliferation, in order to understand the expansion and intensification of dengue and, most importantly, to chose adequate strategies and methods to monitor and control *Ae. aegypti* populations.

A protective tetravalent vaccine must be available if dengue is to be controlled. There are promising vaccine candidates under development (Trindade et al, 2008) and there is a possibility that this will be achieved in a few years (Whitehead et al. 2007, Edelman R 2007). However, the well established *Ae. aegypti* populations in urban environments will remain as a matter of concern since other arboviruses may be transmitted by this species, such as Chikungunya and West Nile viruses.

In Brazilian cities where sensitive surveillance methods were utilized to monitor *Ae. aegypti* populations, such as Salvador (Morato et al, 2005), Manaus (Ríos-Velasquez et al 2007), Recife (Regis et al 2008), the presence of this mosquito species was detected in 60% to 90% of the houses. At the same time, the House Index (HI), used by the National Program for Dengue Control, showed for these places negative or low values, seldom more than 10%. The lack of sensitivity of entomological indices, based on the visualization of immature stages, like the HI and the Breteau Index, is attributed to: a) the strong sensitivity and responsiveness of aedine pre-imaginal forms to water surface vibrations (Slwey et al 1987), and their ability to make deep and long-duration dive and to perform bottom-resting (Romoser and Lucas, 1999), thus making their visualization difficult; b) the information generated is the percentage of house/containers where the presence of one (or more) larva/pupa is observed during a visit that lasts a few minutes and takes place at every two months.

Despite its lack of sensitivity for detecting the presence of *Aedes*, and the fact that it does not quantify population size or density, this method proposed eight decades ago (Connor & Monroe, 1923) is the only entomological surveillance tool used in most programs.

The complexities of the new urban realities point out new targets to the control of mosquito born diseases, such as dengue fever. To increase the capacity of the health sector for controlling transmissible diseases, new tools for epidemiological practice and surveillance

must be developed. The SMCP-*Aedes* – System for Monitoring and Control of the Population of *Aedes aegypti* - is a contribution in this direction.

Dengue vector surveillance is important to determine when and where to apply the control actions. An entomological surveillance system able to generate quantitative information on mosquito population densities and helping to predict explosive population growth is essential for launching actions to prevent epidemics to occur. The importance of the entomologic surveillance as a base for dengue transmission control has been recently highlighted by Ooi et al (2006).

THE SMCP-*Aedes*, A SPATIAL SURVEILLANCE SYSTEM

The main SMCP-*Aedes* concept is to integrate biological aspects of the vector with human population and environmental data through building an entomological surveillance system based on geographical information, as a support for dengue epidemiological surveillance and control at intra-urban scales. To do so the SMCP-*Aedes* makes intensive use of open spatial technology solutions. It is based on open protocols and open standards, an intensive use of web and open software for GIS and Database Management working-up information stored in a spatio-temporal database from where they can be explored through quantitative methods. The *Aedes* population in a certain urban area can be estimated and its spatio-temporal distribution visualized, based on continuous eggs collection. Control interventions can be planned and intensified on the basis of spatial analysis carried on by the SCMP-*Aedes*. These actions to be taken by the local health services can be agreed with the local community. In this sense, the SMCP-*Aedes* can be instrumental for helping the building-up of a permanent social mobilization towards a more effective vector control.

The sample design was conducted supposing to overlap a grid with 400 cells with 50m x 50m each one, for a territory of 1 Km². Depending on the household density in the

area the sample size (number of cells) was calculated by fitting a logistic function (Fig. 1) taking into account the following parameters: 10 cells for very low densities (i.e. 100 or less households per Km²); 100 cells for intermediate density (i.e. almost 3.000 households per Km²) as adopted in a previous experiment (Regis et al 2008); and an upper boundary of 200 (50% of the total) cells for very high densities.

$$Y = \frac{1}{\frac{1}{u} + b_0 \times b_1^X} : \text{ where } Y \text{ is the number of cells to be sampled which will receive one}$$

ovitrap; u is the upper boundary; X is the mean number of households per Km²; and b_0 and b_1 are the parameters estimated by the model.

Each trap was allocated inside the cell space and the coordinates of the location was taken using a Global Positioning System (GPS) and recorded at an integrated geographic information system.

A sensitive, safe and low cost tool is suitable for continuous egg-collection at a number of fixed sampling stations in order to monitor reproductively active populations. This can be accomplished with an ovitrap model consisting of a black plastic cup baited with 2 L of grass infusion (or water supplemented with alfalfa pellets) treated with a Bti-based product and containing three 5 x 15 cm wooden paddles, as oviposition substrates (Regis et al 2008). Besides being a potent, safe and long lasting larvicide specially under shade (Regis et al 2001, Araújo et al 2007), Bti seems to act as an oviposition stimulant (Santos et al 2003, Stoops 2005) conferring increased efficacy to the trap, which can safely remain in the field for up to two months showing an egg-collection capacity of over 7,000 eggs/trap-month (Regis et al 2008). An egg counting computer-assisted system developed by A.F.R. Araújo and collaborators from the Informatic Center, Federal University of Pernambuco, Brazil, allows operational advantages over the optical microscopic counting system. It is less time

consuming, more comfortable for technicians and data records are directly transferred to a geographical database.

THE SMCP-AEDES CONCEPTUAL ARCHITECTURE AND OPEN TECHNOLOGIES

The decision on supporting technology for the SMCP-*Aedes* is based around open software and open standards for geospatial data. It goes in line with our previous work in Recife (Regis et al., 2008) and with the recent works linked with health information systems that have taken the same lines (QianYi et al., 2008; Moreno-Sanches et al., 2007). The database logic model has been based upon the Recife-SAUDAVEL-GED. For the first version of the SMCP-*Aedes* we have taken the open MySQL Database Server as the repository for the geographic entomological database. The MySQL DBMS software is embedded by the open TerraLib GIS library (www.dpi.inpe.br/terralib) allowing for the necessary spatio-temporal models for storing, retrieving and analyzing the data. The eggs counts are recorded on the database using a web-based interface that has been developed by the SAUDAVEL network (Monteiro et al. 2005) at the National Institute for Spatial Research-INPE and at CPqAM/Fiocruz and adapted for the present system. The database contains other data as demographic data at the census tract scale, satellite images of the region and all mapping information available for the sites where the ovitraps have been installed. We have adopted TerraView (www.dpi.inpe.br/terraview), an open GIS application that runs on desktop computers, as a browsing tool for this geographic database which provides quick summaries for some spatio-temporal queries in the stored data, and an integrated environment for visualization of data, remote sensing satellite imagery, pictures of the traps and census tract data.

Within the SMCP-*Aedes* system, assessments and decision making are based on reports systematically and automatically produced incorporating the latest available data. Typically, such outputs make usage of statistical analysis and modeling tools beyond the GIS

capabilities and operational scripts are written using the R statistical computing language (www.r-project.org), an open source environment for statistical analysis (Bonat et al. 2008). These scripts can be modular, implementing requested analysis and report formats and run automatically on the data base server by making use of the open software aRT (leg.ufpr.br/aRT). aRT is a software binder in the format of an R package, integrating the TerraLib GIS library with R. The specific routines for statistical, and in particular spatial and/or temporal analysis, are implemented by another R package, RDengue, tailored to the specific needs of the project and designed to release automatic analysis of the data stored into the SMCP-*Aedes* geographical database (Dallazuanna et al 2008). Fig 2 displays a schematic view of this architecture and detailed schemes, and some example outputs can be seen in Figure 3.

INTEGRATED VECTOR CONTROL BASED ON ENTOMOLOGICAL SURVEILLANCE

The National Program of dengue control, PNCD, established in Brazil in 2002, is implemented by the Ministry of Health in partnership with the 27 states and 5,560 municipalities. The PNCD foresee the implementation of vector control activities, among other components. The vector control is based essentially on the work of health workers carrying out registration of houses and the use of temephos as larvicide. In the last 12 years large amounts of organophosphates have been routinely applied in cycles of two months, with high operational cost. Other insecticides (organophosphates and pyrethroids) are used as adulticides. The application of larvicides and adulticides hardly will reach the control pressure sufficient to reduce populations of this species due to known behavioral and biological characteristics, peculiar to *Ae. aegypti*, in addition to the fact that mosquito populations develop inevitably, resistance to these compounds, as has been detected in the country, since 2000.

The vector control measures proposed here are based on the temporal and spatial distribution of reproductively active mosquito populations, as estimated from eggs collected in the sentinel ovitraps. The aim is to prevent mosquito population outbreaks associated with the seasonal increase of dengue cases by intensifying control measures in specific places at specific times. In Recife, it was observed that the first rainfall after the dry season, associated with the season's high temperatures promoting a massive mosquito egg hatching, resulted in very high mosquito densities from January to August, matching the dengue season. At the same time, entomological spatial analysis allowed the identification of areas where mosquito populations were persistently most concentrated (Regis et al 2008).

The strategy is to combine larval source reduction by eliminating, properly covering or applying Bti as a larvicide in potential larval habitats, with mass collection/destruction of eggs, through a pact between public health services and the local community. In this strategy, the use of ovitraps is essential to avoid gravid females to disperse in the search for oviposition sites. The public must be aware that the lower the amount of available breeding sites in a house, the higher the amount of eggs that will be laid in an ovitrap. Another important aspect is that the oviposition substrate used in these ovitraps (cotton fabric, according to Lenhard 2005) allows the visualization of *Aedes* eggs by the householders, and it is easier to be incinerated (for egg destruction) than wooden substrates. For security reasons the Government should take responsibility for egg incineration and supply of larvicides based on Bti to be used in the ovitraps.

Besides these actions, aiming to avoid a general mosquito population growth, the strategy also includes control interventions targeting the hotspots, focusing on mechanical elimination of adult mosquitoes from houses located in these areas. The use of aspirators for indoor collection of adult mosquitoes has several advantages over the use of organophosphorous or pyrethroids as adulticides: it does not offer any health risk for humans

and domestic animals; the method is efficient and effective since mosquitoes are caught directly from their resting places within the house; householders are able to see the mosquito capture; results are measurable. It is expected that the vector population is impacted by indoors mosquito captures repeated fortnightly for two consecutive months within houses containing two or three ovitraps for egg collection-destruction and where potential breeding sites are properly managed.

FINAL REMARKS

Built on an interdisciplinary approach, the SMCP-*Aedes* was developed and assessed based on a longitudinal study conducted at Recife-PE over a 4 years period (Regis et al 2008). It has now been putting into action as a pilot program in two cities of the Pernambuco state, Brazil, in a joint effort involving the state health authority, the local health services, research institutions (Fiocruz and INPE) and the SAUDAVEL network (Monteiro et al 2005). The system is designed to generate automatic analysis and reports and uses open software produced in Brazil namely the TerraLib/TerraView, aRT and RDengue, combined with a R software for statistical computing and a database management system. The following-up of the outcomes of the program in the pilot cities of Santa Cruz do Capibaribe-PE and Ipojuca-PE is expected to demonstrate how open spatial information technology can contribute to establish an entomological surveillance strongly based on open spatial technology solutions with low cost and high efficiency.

RESUMO

Associadas a fatores bióticos, climáticos e culturais que favorecem a proliferação do *Aedes aegypti* em ecossistemas urbanos, adaptações evolutivas à hematofagia freqüente e quase exclusiva em humanos e à colonização de tipos infinitos de habitats, fazem deste mosquito um disseminador extremamente eficiente de patógenos ao homem. Estudos utilizando

métodos sensíveis de monitoramento demonstram que os métodos de vigilância usados no programa brasileiro não revelam as elevadas intensidades da infestação das cidades por este vetor. Para ampliar a capacidade do setor saúde novos instrumentos são necessários à prática da vigilância, incorporando aspectos do vetor, do lugar e das pessoas do lugar. Apresentamos aqui o SMCP-*Aedes* - Sistema de Monitoramento e Controle Populacional do *Ae. aegypti*, cuja meta é a instrumentalização da vigilância entomológica como base para a vigilância epidemiológica da dengue. Para isso ele se apóia em tecnologias da informação espacial baseadas no uso intensivo da web e de software livre para coletar, armazenar, analisar e disseminar informações relativas à distribuição espaço-temporal da densidade estimada para a população do *Aedes*, com base em amostras obtidas continuamente com ovitrampas. Intervenções de controle planejadas e intensificadas onde e quando indicado pela vigilância entomológica, são pactuadas com os habitantes, apoiando-se na mobilização social permanente.

Palavras-chave: controle do vetor, dengue, vigilância entomológica, ovitrampas, SIG.

ACKNOWLEDGMENTS: This work was supported by CNPq, Fiocruz-PDTSP/Rede Dengue, MS-Decit/UNESCO, and FACEPE.

REFERENCES

- ARAÚJO AP, MELO-SANTOS MAV, CARLOS SO, RIOS EMM AND REGIS L. 2007. Evaluation of an experimental product based on *Bacillus thuringiensis* sorovar. *israelensis* against *Aedes aegypti* larvae (Diptera: Culicidae). Biol Control 41: 339-347.
- BONAT WR, DALLAZUANNA HS, RIBEIRO JR PJ. 2008. Investigando fatores associados a contagens de ovos de *Aedes aegypti* coletados em ovitrampas em Recife, PE. Relatório Técnico, Laboratório de Estatística e Geoinformação, UFPR.
- CONNOR ME AND MONROE, WM. 1923. *Stegomyia* indices and their value in yellow fever control. Am J Trop Med Hyg 3: 9-19.
- DALLAZUANNA HS, BONAT WR, RIBEIRO JR PJ. 2008. RDengue: pacote de apoio à análise para monitoramento de ovos do *Aedes aegypti*. Versão 1.0

- EDELMAN R. 2007. Dengue Vaccines Approach the Finish Line. *Clin Infect Dis* 45: S56–S60.
- HARRINGTON LC, EDMAN JD AND SCOTT TW. 2001. Why do female *Aedes aegypti* (Diptera: Culicidae) feed preferentially and frequently on human blood? *J Med Entomol* 38: 411-422.
- HARRINGTON LC, OISEVERMEYLEN F, JONES JJ, WEE SK, SITHIPRASASNA R, EDMAN JD AND SCOTT TW. 2008. Age-Dependent Survival of the Dengue Vector *Aedes aegypti* (Diptera: Culicidae) Demonstrated by simultaneous Release–Recapture of Different Age Cohorts. *J Med Entomol* 45: 307-313.
- KOENRAADT CJ AND MHARRINGTON LC. 2008. Flushing Effect of Rain on Container-Inhabiting Mosquitoes *Aedes aegypti* and *Culex pipiens* (Diptera: Culicidae) *J Med Entomol* 45: 28-35
- LENHART AE, WALLE M, CEDILLO H AND KROEGER A 2005. Building a better ovitrap for detecting *Aedes aegypti* oviposition. *Acta Trop* 96: 56-59.
- MONTEIRO AMV, CARVALHO MS, ASSUNÇÃO R, SOUZA WV, RIBEIRO JR PJ, DAVIS JR C, REGIS L, the SAUDAVEL Project Team. 2005. SAUDAVEL: Bridging the Gap between Research and Services in Public Health Operational Programs by Multi-Institutional Networking Development and Use of Spatial Information Technology Innovative Tools. [cited 2008 Jan 10]. Available from: <http://www.dpi.inpe.br/saudavel/publicacoes.html>.
- MORATO VCG, TEIXEIRA MG, GOMES AC, BERGAMASCHI DP AND BARRETO M 2005. Infestation of *Aedes aegypti* estimated by oviposition traps in Brazil. *Rev Saude Pub* 39: 553-558.
- MORENO-SANCHEZ R, ANDERSON G, CRUZ J AND HAYDEN M. 2007 The potential for the use of Open Source Software and Open Specifications in creating Web-based cross-border health spatial information systems. *Int J Geogr Inf Sci* 21 (10): 1135-1163.
- OOI E-E, GOH K-T AND GUBLER DJ. 2006 Dengue Prevention and 35 Years of Vector Control in Singapore. *Emerg Infect* 12 (6): 887-893.
- PONLLAWAT A AND HARRINGTON L. 2005. Blood feeding patterns of *Aedes aegypti* and *Aedes albopictus* in Thailand. *J Med Entomol* 42: 844-849.
- QIAN YI ET AL. 2008. Integrating open-source technologies to build low-cost information systems for improved access to public health data. *Int J Health Geogr* 7: 29.
- REGIS L, SILVA-FILHA MH, NIELSEN-LEROUX C AND CHARLES JF 2001. Bacteriological larvicides of dipteran disease vectors. *Trends Parasitol* 17: 377-380.

- REGIS ET AL 2008 Developing new approaches for detecting and preventing *Aedes aegypti* population outbreaks: basis for surveillance, alert and control system. Mem Inst Oswaldo Cruz 103 (1): 50-59.
- RÍOS-VELASQUEZ CM, CODEÇO CT, HONÓRIO NA, SABROSA PS, MORESCO M, CUNHA ICL, LEVINO A, TOLEDO LM AND LUZ SLB. 2007. Distributions of dengue vectors in neighborhoods with different urbanization types of Manaus, state of Amazonas, Brazil. Mem Inst Oswaldo Cruz 102: 617-623.
- ROMOSER WS AND LUCAS EA. 1999. Buoyancy and diving behavior in mosquito pupae. J Am Mosq Contr Assoc 15: 194-199.
- SANTOS SRA, MELO-SANTOS MAV, REGIS L AND ALBUQUERQUE CMR. 2003. Field evaluation of ovitraps consociated with grass infusion and *Bacillus thuringiensis* var. *israelensis* to determine oviposition rates. Dengue Bull 27: 156-162.
- STOOPS CA 2005. Influence of *Bacillus thuringiensis* var. *israelensis* on oviposition of *Aedes albopictus* (Skuse). J Vector Ecol 30: 41-44.
- SCOTT TW, CHOW E, STRICKMAN D, KITTAYAPONG P, WIRTZ RA AND J.D. EDMAN. 1993. Blood feeding patterns of *Aedes aegypti* collected in a rural Thai village. J Med Entomol 30: 922-927.
- SCOTT TW, AMERASINGHE PH, MORRISON AC, LORENZ LH, CLARK GG, STRICKMAN D, KITTAYAPONG P AND EDMAN JD. 2000. Longitudinal studies of *Aedes aegypti* (L.) (Diptera: Culicidae) in Thailand and Puerto Rico: Blood feeding frequency. J Med Entomol 37: 89-101.
- SCOTT TW AND MORRISON AC. 2003. *Aedes aegypti* density and the risk of dengue virus transmission. In: *Ecological aspects for application of genetically modified mosquitoes*. eds. W. Takken and T.W. Scott. FRONTIS, Dordrecht, The Netherlands.pp: 187-206.
- SHUEY JA, BUCCI AJ AND ROMOSER WS. 1987. A behavioral mechanism for resting site selection by pupae in three mosquito species. J Am.Mosq Control Assoc 3: 65-69.
- TRINDADE GF, MARCHEVSKY RS, DE FILLIPIS AMB, NOGUEIRA RMR, BONALDO MC, ACERO PC, CARIDE E, FREIRE MS AND GALLER R. 2008. Limited replication of yellow fever 17DD and 17D-Dengue recombinant viruses in rhesus monkeys. An Acad Bras Cienc 80: 311-321.
- WHITEHEAD SS, BLANEY JE, DURBIN AP AND MURPHY BR. 2007. Prospects for a dengue virus vaccine. Nat Rev Microbiol 5: 518-528.

FIGURE CAPTIONS

Figure 1. Logistic Function describing the number of cells to be sampled according to the household density.

Figure 2. A General Conceptual Scheme for the Computational Architecture of the SMCP-*Aedes* including the open set of spatial technologies chosen for implementation of the system. It involves open GIS library TerraLib, the open GIS application TerraView, the open database management system MySQL and the aRT- an Application Programming Interface for binding R to TerraLib running the RDengue statistical analysis package.

Figure 3. (a) A simplified scheme showing the interaction amongst an open DBMS system, the open TerraLib GIS library and open statistical computing environment R; (b) A schematic view on how the open technologies work together to promote spatial analysis and visualization of the entomological and geographical data; (c) Example of analysis carried out by the RDengue to a specific ovitrap showing its localization and statistical summaries relating average egg counting on that trap against its neighbors over time; (d) Another example of RDengue possibilities showing a *hotspot* map for a specific site displaying the estimated vector density based on the ovitraps counting over a certain period.




Fig 1

SMCP-*Aedes* Conceptual View and Supporting Technologies




Figure 2




Figure 3